

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number
WO 2004/053140 A2

(51) International Patent Classification⁷: C12Q
(21) International Application Number:
PCT/IN2003/000389

(22) International Filing Date:
12 December 2003 (12.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
925/MAS/2002 12 December 2002 (12.12.2002) IN

(71) Applicant (for all designated States except US): JAWA-
HARLAL NEHRU CENTRE FOR ADVANCED SCI-
ENTIFIC RESEARCH [IN/IN]; Jakkur Post, Bangalore
560 064, Karnataka (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KUNDU, Tapas,
Kumar [IN/IN]; No. JJ-D1 type Quarters, Jawaharlal
Nehru Centre for Advanced Scientific Research, Jakkur
Campus, Jakkur Post, Bangalore 560 064, Karnataka (IN).
BALASUBRAMANYAM, Karanam [IN/IN]; No. 145,
Padmalaya, 2nd H Main, 11th Block, II Stage, Nagarbhavi,
Bangalore 560 072, Karnataka (IN). SWAMINATHAN,
Venkatesh [IN/IN]; "Sanskar", Flat No. 1A, 8 Lake
Range, Kolkata 700 026, West Bengal (IN).

(74) Agent: VAIDYANATHAN, Alamelu; 451, 2nd Cross, 3rd
Block, 3rd Stage, Basaveshwaranagar, Bangalore 560 079,
Karnataka (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT,
TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent
(BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG)


— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for the following desig-
nations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY,
BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC,
EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW,
ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ,
TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY,
CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: MODULATORS (INHIBITORS/ACTIVATORS) OF HISTONE ACETYLTRANSFERASES



(I)



(II)

(57) Abstract: The invention provides the use of certain benzoic acid and benzamide compounds as modulators of enzymes histone acetyltransferases, which are involved in gene expression and cancer and also use of such compounds in the treatment of diseases due to defects in gene regulation predominantly cancer.



— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Modulators (inhibitors/activators) of histone acetyltransferases

Field of invention:

This invention relates to the field of novel anticancer agents for therapeutic application in human medicine.

5 Background:

The eukaryotic genome is organized as a highly complex nucleoprotein structure called chromatin, the unit of which is the nucleosome. The nucleosome is composed of two copies each of four different histones, H3, H2B, H2A, and H4, constituting a scaffold, which is wrapped around by 146 base pairs of DNA. Therefore, for any
10 process that requires access to the DNA (*e.g.* transcription, replication, recombination, and repair), the chromatin needs to be opened by the remodeling systems. There are two different biochemical processes to modify chromatin structure, namely the covalent modifications of histone tails and the ATP-dependent chromatin remodeling. Among the several covalent modifications of histones known, the reversible
15 acetylation of key lysine residues in histones holds a pivotal position in transcriptional regulation. Acetylation of histones is a distinctive feature of the transcriptionally active genes, whereas deacetylation indicates the repressed state of a gene,. A balance between the acetylation and deacetylation states of histones regulates transcription. Dysfunction of the enzymes involved in these events, the histone acetyltransferases
20 (HATs) and histone deacetylases (HDACs), is often associated with the manifestation of cancer. These enzymes thus become potential new targets for antineoplastic therapy.

A wide repertoire of transcriptional co-activator proteins is now recognized to possess histone acetyltransferase activity. These include p300/CBP-associated factor (PCAF),
25 which is similar to GCN5, nuclear hormone receptor cofactors such as steroid receptor cofactor 1 (SRC1) and activator of thyroid and retinoid receptor (ACTR) and the multifunctional p300/CBP. The p300/CBP is a global transcriptional coactivator, which plays a critical role in a variety of cellular process including cell cycle control, differentiation, and apoptosis. Mutations in p300/CBP are associated with different
30 human cancers and other human diseases. It is one of the most potent histone

acetyltransferases, which can acetylate all four-core histones within nucleosomes as well as free histone forms. The HAT activity of p300 is regulated by several other factors. For example, the viral oncoprotein E1A binds to p300 and inhibits its activity, whereas phosphorylation of CBP by cyclin E/Cdk2 kinase activates its HAT activity.

- 5 During the process of transcription, p300 is recruited on to the chromatin template through the direct interaction with the activator and enhances the transcription by acetylation of promoter proximal nucleosomal histones.

- Although significant progress has been made in the field of histone deacetylase inhibitors as antineoplastic therapeutics, and some of the compounds are already in
10 human trials the reports of HAT inhibitors/activators are scanty. Prior to the molecular characterization of HAT enzymes, several polyamine-CoA conjugates were found to block HAT activity in cell extracts. However, the target enzyme(s) for these conjugates was not known. Recently, two peptide-CoA conjugates, namely Lysyl CoA (Lys-CoA) and H3-CoA-20, were synthesized that specifically inhibit the HAT
15 activity of p300 and PCAF, respectively. Thus, there is an urgent medical need to identify new drugs having modulating activity(inhibitors/activators) towards histone acetyltransferases which can be used to treat diseases in which histone acetyltransferase play an important role treatment of diseases due to defects in gene regulation predominantly cancer.

- 20 In accordance with the present invention, there is therefore provided the use of a compound of general formula I for activator molecules of histone acetyltransferases and the use of a compound of general formula II for inhibitor molecules of histone acetyltransferases.

Description of the Invention:

- 25 Cashew nut shell liquid (CNSL) possessed inhibitory activity towards p300. The systematic bio-activity guided fractionation of CNSL yielded unsaturated anacardic acids mixture, namely, the 8'Z-monoene, the 8'Z, 11'Z-diene, and the 8'Z, 11'Z, 14'Z-triene, which are the chief constituents (~75%) of cashew nutshell liquid (18), having maximum HAT inhibitory activity. The hydrogenation of unsaturated

30

anacardic acids mixture yielded a single compound, Anacardic acid (2-hydroxy-6-pentadecylbenzoic acid) showing an equally potent inhibitory activity towards p300. This data indicated that absence of unsaturation in anacardic acid did not alter its HAT inhibitory property. Neither salicylic acid nor benzoic acid shown any
5 inhibitory activity against histone acetyltransferases.

The rate of the acetylation reaction at different concentrations of the inhibitors (and in its absence) was recorded with increasing concentrations of [^3H]-acetyl CoA and a constant amount of core histones. The double reciprocal plot for each inhibitor concentration and in its absence ($1/\text{c.p.m}$ vs. $1/[\text{Acetyl CoA}]$) was plotted as shown in
10 Figure 2E. The results suggest that anacardic acid is a non-competitive type of p300-HAT inhibitor.

The acidic group on the both salicylic acid and anacardic acid was modified to respective different amide derivatives using substituted anilides. Most of these compounds, with different anilide moieties on anacardic acid and salicylic acid, when
15 tested *in vitro* HAT assay (filter binding), surprisingly showed an enhancement in the p300 HAT activity, while keeping the PCAF HAT activity mostly unperturbed. The interchanging of substitution pattern on the ring B in formula I was found to be affecting activation profile of molecule.

Deacetylation of the core histones in the presence or absence of the HAT activating
20 compounds, at 100 μM or 500 μM , shows no difference in the deacetylase enzyme activity indicating no affect on deacetylation activity and the HAT enzyme specificity.

The addition of increasing concentration of either inhibitors or activators does not produce any variation in the transcript levels as compared to the DMSO control indicating that the compounds do not affect any component of the basal transcription
25 machinery.

The template pG₅ML-array (8) was assembled into chromatin using the NAP1 mediated assembly method (Experimental Procedures). Addition of activators to the HAT-dependent transcription reaction along with the p300 and acetyl CoA after

allowing for 30 min of acetylation either in the presence or absence of the compound. Under these conditions we found the addition of DMSO produced a slight drop in the transcript levels while the addition of activators enhanced the levels of transcription 1.6 fold over the DMSO control. Thus this result indicates that CTPB specifically
5 enhances the HAT activity of p300, a function that is reflected even at the transcriptional level. In order to explain the 1.6-fold increase in transcription levels, in contrast to the ~5-fold increase in the histone acetylation levels; Inhibitors did not affect the transcription from the DNA template, but the HAT-dependent transcription from chromatin template was inhibited by addition of inhibitors at 10µM
10 concentration.

Methods:

A: Chemical Methods

The cashew nut shell liquid (CNSL) is also known as cashew nut shell oil. CNSL is a dark brown viscous liquid reported to be 15-20% by weight of the unshelled nut in
15 Africa, 25-30% by weight in India and ca. 25% overall. CNSL contains 90% anacardic acid and 10% cardol.

Methods reported to extract CNSL from cashew shells are:

1. Roasting nuts and collecting expelled liquid (Indian native method, yield 50%).
2. Extract with hot CNSL without charring the kernels (yield 85-90%).
- 20 3. Super heated steam treatment and collect condensate (used to improve yields of 2).
4. Solvent extraction with hexane leading to more percentage of anacardic acid.
5. Supercritical fluid extraction.

CNSL can also be extracted from cashew nuts that are soaked in water or humidified
25 in piles and then held in a humid atmosphere so that the shell has set moisture content from 15-45% depending on the methods. CNSL is used in the manufacture of brake linings, industrial belting and clutches, reinforcing synthetic rubber, for oil and acid resistance, in lacquers, in electrical insulation material, as a metal anti-corrosive

material, for waterproofing and as an adhesive. The use of CNSL in varnishes, lacquers, paints and brake linings requires distillation and further refinement.

Anacardic acid (6-pentadecylsalicylic acid), a major component of cashew nut shell liquid (CNSL), is obtained by solvent extraction of cashew nut shells. It exists as a heterogeneous mixture of monoenes, dienes, and trienes. More specifically Cold processed CNSL was purchased from commercial source. The Anacardic acid present in the CNSL was purified as calcium anacardate by adding Calcium hydroxide to CNSL dissolved in isopropyl alcohol. The pure calcium salt of anacardic acid was dried and treated with 1N HCl to release free anacardic acid ene mixture (containing $n=0,2,4,6$) (General food corporation (Rye, NY) Indian patent, 34671,1946). The ene mixture obtained by above method was hydrogenated in ethylacetate for 4hrs over 10% palladium-carbon using a Parr hydrogenator. The catalyst was filtered off and the solvent evaporated in vacuo to yield saturated anacardic acid. The alkylation with dimethyl and diethyl sulphates using potassium carbonate gave the dialkylated derivative. Di isopropyl anacardic acid was obtained by using isopropyl bromide in presence of K_2CO_3 with phase transfer catalyst in M1BK for 36 hrs.

Dialkylated anacardic acids were treated with potassium tertiary butoxide in DMSO to yield respective O-alkyl anacardic acids. The O-alkyl anacardic acids on treatment with thionyl chloride in the presence of a catalytic amount of DMF yield corresponding O-alkyl anacardic acid chlorides. The resultant acid chlorides condensed with different substituted anilines yielded respective benzamide derivatives.

All compounds were characterized by using FABMS. TLC was done using precoated silica gel GF₂₅₄ plates (Merck, Darmstadt, Germany) with hexane: Ethyl acetate (7:3) as the developing solvent and visualized after spraying with vanillin sulphuric acid reagent.

The following are preparation of starting materials used for synthesis of activator molecules and inhibitor molecules.

Isolation of Anacardic Acid from CNSL. Commercially available solvent-extracted CNSL (1Kg) was dissolved in 5% aqueous methanol (6.2L), and calcium hydroxide (510 g) was added in portions under stirring. After complete addition of calcium hydroxide, the temperature of the reaction mixture was raised to 54 °C and stirring
5 was continued for 4h. The supernatant solution was monitored by TLC for the absence of anacardic acid. After completion of the reaction, the precipitated calcium anacardate was filtered and washed thoroughly with methanol (2.5L), and the cake was dried under vacuum at 45-54 °C for 3h (dry weight 1.08 Kg). The filtrate was preserved for subsequent isolation of cardol and cardanol. Calcium anacardate
10 (1.08Kg) was suspended in distilled water (4.5 L) and 11 M HCl (600 mL) was added and stirred for 1.5 h. The resultant solution was extracted with ethyl acetate (2 × 2 L). The combined organic layer was washed with distilled water (2 × 2L), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to yield 600 g of mixture of anacardic acid (monoene, diene, and triene).

15 **Preparation of Saturated Anacardic Acid .** To a solution of an ene mixture of anacardic acid (500 g, 1300 mmol) in methanol (2.0 L) was added 5% palladium-carbon (25 g), and then hydrogen gas was passed through the solution at 2.5 kg/cm² until consumption of hydrogen gas ceased (4-5 h). The reaction mass was filtered over a Celite bed and washed with methanol (100 mL). The filtrate was concentrated under
20 reduced pressure to give an off-white solid (477 g), which on recrystallization from hexane yielded 450g .

Preparation of Methyl 2-methoxy-6-pentadecylbenzoate. To a stirred solution of anacardic acid mentioned above (50 g, 143 mmol) in acetone (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). Dimethyl sulfate (44.25
25 g, 290 mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic
30 layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Methyl 2-methoxy-6-pentadecylbenzoate (45 g, 80%).

Preparation of 2-Methoxy-6-pentadecylbenzoic Acid. To a stirred solution of Methyl 2-methoxy-6-pentadecylbenzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Methoxy-6-pentadecylbenzoic Acid (7.6 g, 80%).

Preparation of 2-Methoxy-6-pentadecylbenzoyl Chloride. To a stirred solution of 2-Methoxy-6-pentadecylbenzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Methoxy-6-pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides.

Preparation of Ethyl 2-Ethoxy-6-pentadecylbenzoate. To a stirred solution of anacardic acid mentioned above (50 g, 143 mmol) in acetone (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). Diethyl sulfate (44.25 g, 290 mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Ethyl 2-Ethoxy-6-pentadecylbenzoate (45 g, 80%).

Preparation of 2-Ethoxy-6-pentadecylbenzoic Acid. To a stirred solution of Ethyl 2-Ethoxy-6-pentadecylbenzoate (20 g, 240 mmol) in dimethyl sulfoxide (80 mL) was

added potassium *tert*-butoxide (20 g, 890 mmol) in portions. The solution was heated to 68 °C on a water bath for 3 h, the reaction was monitored by TLC using a hexane-ethyl acetate (9:1) solvent system. The reaction mass was cooled to 15°C, poured into ice water, and then acidified with 10% dilute hydrochloric acid. The precipitated solid
5 was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in petroleum ether (50 mL) to yield an off-white solid of 2-Ethoxy-6-pentadecylbenzoic Acid (15 g, 80%).

Preparation of 2-Ethoxy-6-pentadecylbenzoyl Chloride. To a stirred solution of 2-Ethoxy-6-pentadecylbenzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added
10 thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Ethoxy-6-pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides .

Preparation of Propyl 2-propoxy-6-pentadecylbenzoate. To a stirred solution of anacardic acid mentioned above (50 g, 143 mmol) in acetone (300 mL) was added
15 anhydrous powdered potassium carbonate (80 g, 580 mmol). *n*-propyl iodide (44.25 g, 290 mmol) was added in portions for about 24hrs at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath
20 and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Propyl 2-propoxy-6-pentadecylbenzoate (45
25 g, 80%).

Preparation of 2-Propoxy-6-pentadecylbenzoic Acid. To a stirred solution of Propyl 2-propoxy-6-pentadecylbenzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution
30 was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute

hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Propoxy-6-pentadecylbenzoic Acid (7.6 g, 80%).

- 5 **Preparation of 2-Propoxy-6-pentadecylbenzoyl Chloride.** To a stirred solution of 2-Ethoxy-6-pentadecylbenzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Propoxy-6-
10 pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides

- Preparation of Isopropyl 2-isopropoxy-6-pentadecylbenzoate.** To a stirred solution of anacardic acid mentioned above (50 g, 143 mmol) in acetone (300 mL) was added
15 anhydrous powdered potassium carbonate (80 g, 580 mmol). isopropoxyiodide (44.25 g, 290 mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the
20 reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Isopropyl 2-isopropoxy-6-pentadecylbenzoate (45 g, 80%).

- 25 **Preparation of 2-Isopropoxy-6-pentadecylbenzoic Acid.** To a stirred solution of Isopropyl 2-isopropoxy-6-pentadecylbenzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The
30 reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly

with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Isopropoxy-6-pentadecylbenzoic Acid (7.6 g, 80%).

Preparation of 2-Isopropoxy-6-pentadecylbenzoyl Chloride. To a stirred solution of 2-Isopropoxy-6-pentadecylbenzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Isopropoxy-6-pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for condensation with anilides .

10 **Preparation of Methyl 2-methoxy-benzoate.** To a stirred solution of salicylic acid (50 g, 143 mmol) in acetone (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). Dimethyl sulfate (44.25 g, 290 mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Methyl 2-methoxy-benzoate (45 g, 80%).

20 **Preparation of 2-Methoxy-benzoic Acid.** To a stirred solution of V (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Methoxy-benzoic Acid (7.6 g, 80%).

Preparation of 2-Methoxy-benzoyl Chloride. To a stirred solution of 2-Methoxy-benzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to

reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Methoxy-benzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides.

- 5 **Preparation of Ethyl 2-Ethoxy-benzoate.** To a stirred solution of Salicylic acid (50 g, 143 mmol) in acetone (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). Diethyl sulfate (44.25 g, 290 mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Ethyl 2-Ethoxy-benzoate (45 g, 80%).

- 15 **Preparation of 2-Ethoxy-benzoic Acid.** To a stirred solution of Ethyl 2-Ethoxy-benzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Ethoxy-benzoic Acid (7.6 g, 80%).

- 25 **Preparation of 2-Ethoxy-benzoyl Chloride.** To a stirred solution of 2-Ethoxy-6-pentadecylbenzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Ethoxy-benzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides.

Preparation of Propyl 2-propoxy-6-pentadecylbenzoate. To a stirred solution of Salicylic acid mentioned above (50 g, 143 mmol) in ACETONE (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). n-propyl iodide (44.25 g, 290 mmol) was added in portions for about 24hrs at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Propyl 2-propoxy-benzoate (45 g, 80%).

Preparation of 2-Propoxy-benzoic Acid. To a stirred solution of Propyl 2-propoxy-benzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Propoxy-benzoic Acid (7.6 g, 80%).

Preparation of 2-Propoxy-benzoyl Chloride. To a stirred solution of 2-Propoxy-benzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Propoxy-6-pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for the condensation with anilides .

Preparation of Isopropyl 2-isopropoxy-benzoate. To a stirred solution of anacardic acid mentioned above (50 g, 143 mmol) in acetone (300 mL) was added anhydrous powdered potassium carbonate (80 g, 580 mmol). isopropyl iodide (44.25 g, 290

30

mmol) was added in portions for about 10 min at room temperature. After the addition was complete, the solution was heated to reflux temperature on a water bath and maintained for 3 h. The solution was cooled to room temperature and then concentrated under reduced pressure. Distilled water (200 mL) was added to the reaction mixture, which was then extracted with ethyl acetate (200 mL). The organic layer was washed with distilled water (2 × 200 mL), dried over anhydrous sodium sulfate, and concentrated to yield the title Isopropyl 2-isopropoxy-benzoate (45 g, 80%).

Preparation of 2-Isopropoxy-benzoic Acid. To a stirred solution of Isopropyl 2-isopropoxy-benzoate (10 g, 240 mmol) in dimethyl sulfoxide (40 mL) was added potassium *tert*-butoxide (10 g, 890 mmol) in portions. The solution was heated to 70 °C on a water bath for 2 h, and the progress of the reaction was monitored by TLC using a hexane-ethyl acetate (8:2) solvent system. The reaction mass was cooled to 10 °C, poured into ice water, and then acidified with 5% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in hexane (50 mL) to yield an off-white solid of 2-Isopropoxy-benzoic Acid (7.6 g, 80%).

Preparation of 2-Isopropoxy-benzoyl Chloride. To a stirred solution of 2-Isopropoxy-benzoic Acid (6.5 g, 16 mmol) in hexane (60 mL) were added thionyl chloride (2.5 g, 21 mmol) and *N,N*-dimethylformamide (0.5 mL). The reaction mixture was heated to reflux for 1 h. After the reaction was complete, the solvent was evaporated under reduced pressure to yield the desired 2-Isopropoxy-6-pentadecylbenzoyl Chloride, which was redissolved in dichloromethane (50 mL) and used for condensation with following anilides to yield respective benzamides.

25 ***B: Biochemical Methods:***

Purification of Human Core Histones and Recombinant Proteins— Human core histones were purified from HeLa nuclear pellet as described previously (13). The FLAG epitope tagged Human Topoisomerase I, histone deacetylase 1 (HDAC1) and PCAF, were purified from the recombinant baculovirus infected insect cell line, Sf21,

30

- by the immunoaffinity purification using M2-agarose (SIGMA) (14). Full-length p300 was also purified from the recombinant baculovirus infected Sf21 cells as a His₆-tagged protein through the Ni-NTA affinity column (Qiagen) as described previously (13). The His₆-tagged nucleosome assembly protein 1 (NAP1), used for
- 5 the *in vitro* chromatin assembly was purified from *E.coli* cells as previously reported (13) and the FLAG-tagged chimeric activator Gal4-VP16, expressed in *E. coli* and purified by immunoaffinity purification with M2 agarose. Human positive transcriptional coactivator, PC4, was expressed in *E.coli* and purified as described earlier (15).
- 10 The peptide substrate, a 45 residue core histone H3 N-terminal peptide (N-CARTKQTARKSTGGKAPRQLASKAARKSAPSTGGVKKPHRYKPG-C) was synthesized.

- HAT Assay:** Each of the compounds of Examples 1 to 26 were assessed for inhibition / activation of histone acetyltransferase activity of enzymes p300, PCAF and CBP.
- 15 HAT assays were performed as described elsewhere (13). Briefly, indicated amounts of proteins/peptide were incubated in HAT-assay buffer containing 50 mM Tris-HCl, pH 8.0, 10% (v/v) glycerol, 1 mM dithiothreitol, 1 mM phenylmethyl sulfonyl fluoride, 0.1 mM EDTA, pH 8.0, 10 mM sodium butyrate at 30 °C for 10 min in presence or absence of compound followed by the addition of 1 µl of 6.2 Ci/mmol
- 20 [³H]-acetyl Coenzyme A (acetyl-CoA) and were further incubated for another 10 min. The final reaction volume was 30µl. The reaction mixture was then blotted onto P-81 (Whatman) filter papers and radioactive counts were recorded on a Wallac 1409 liquid scintillation counter.

- In order to characterize the inhibition kinetics of anacardic acid, filter-binding assays
- 25 were done using constant amount of HeLa core histones in the presence or absence of AA with increasing concentrations of [³H]-acetyl CoA (see Figure Legends, 2E). To visualize the radiolabeled acetylated histones, the reaction mixtures were resolved on 15% SDS-PAGE and processed for fluorography as described elsewhere (15).

Histone deacetylase assay: Deacetylation assays were performed in the HAT assay buffer without sodium butyrate. 2 µg of core histones were incubated with 20 ng of p300 and 1 µl of 6.2 Ci/mmol [³H]-acetyl CoA for 15 min at 30°C. The activity of p300 was inhibited by incubating the reaction mixture with 10 nM p300-HAT specific inhibitor, Lysyl-CoA (10), for 10 min after which 50 ng of HDAC1 was added, in the presence or absence of the compounds, and incubated further for 45 min. The samples were analysed by flourography.

In Vitro Chromatin assembly: Chromatin template for *in vitro* transcription experiments was assembled and characterized as described earlier (8).

10 **In Vitro Transcription Assay:** Transcription assays were essentially carried out as described elsewhere (8), with minor modifications. The scheme of transcription is enumerated in Figure 5A. Briefly, 30 ng of DNA/equivalent amount of chromatin template was incubated with 30 ng of activator (Gal4-VP16) in a buffer containing 4 mM HEPES (pH 7.8), 20 mM KCl, 2 mM DTT, 0.2 mM PMSF, 10 mM sodium butyrate, 0.1 mg/ml BSA, 2% glycerol (8). The activating compounds were added to
15 the acetylation reaction along with p300 and acetyl-CoA, and incubated for 30 min. at 30°C. This was followed by addition of the p300 specific inhibitor Lysyl CoA (5 µM) to quench the acetylation reaction.

For inhibitors, the HAT p300 was pre-incubated with indicated amounts of inhibitor
20 on ice for 20 min., following which it was added to the acetylation reaction in the transcription assay. For the DNA transcription assays and chromatin transcription inhibition assays, the Lysyl CoA step was omitted. After acetylation, HeLa nuclear extract (5 µl, which contains ~8 mg/ml protein) was added to initiate the pre-initiation complex formation. Transcription reaction was started by the addition of NTP-mix
25 and α-[³²P]-UTP, after the pre-initiation complex formation. The incubation was continued for 40 min at 30°C. Transcription was terminated by the addition of 250 µl stop buffer (20 mM Tris-HCl pH 8.0, 1 mM EDTA, 100 mM NaCl, 1% SDS and 0.025 ng/µl tRNA). The ³²P-radiolabeled transcript was extracted with phenol-chloroform, ethanol precipitated, dried pellet dissolved in loading dye (8 M Urea,

30

0.005% bromophenol blue and xylene cyanol) and analyzed on 5% urea-polyacrylamide gel. Gels were then dried and subjected to autoradiography at -70°C. Quantification of transcription was done by Fuji BAS system. Quantitation of DNA and chromatin transcription data represents three independent experiments.

- 5 The following Examples illustrate the preparation of compounds of formula I using the starting materials described above:

EXAMPLE 1:

2-Ethoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide:

- 2-ethoxy--benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in
10 dichloromethane in presence of triethylamine as acid scavenger to yield 2-Ethoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained
15 was chromatographed over silica gel to afford the desired product.

¹H-NMR:δ6.95-8.10 (7H,aromatic), δ3.98(2H,t,OCH₂), δ1.33(3H,t,Methyl)

EXAMPLE 2:

2-Methoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide:

- 2- Methoxy-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in
20 dichloromethane in presence of triethylamine as acid scavenger to yield 2-Methoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained
25 was chromatographed over silica gel to afford the desired product.

¹H-NMR:δ6.95-8.10 (7H,aromatic), δ3.73(3H,t,Methyl).

EXAMPLE 3:

2-Propoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide:

2-Propoxy-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield 2-Propoxy-N-
5 (4-nitro-3-trifluoromethyl-phenyl)-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

10 ¹H-NMR: δ 6.95-8.10 (7H, aromatic), δ 3.94(2H, t, OCH₂), δ 1.75(2H, t, Methylene) δ 0.96 (3H, t, Methyl).

EXAMPLE 4:

2-Isopropoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide:

2-Isopropoxy-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride
15 in dichloromethane in presence of triethylamine as acid scavenger to yield 2-Isopropoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained
20 was chromatographed over silica gel to afford the desired product.

¹H-NMR: δ 6.95-8.10 (7H, aromatic), δ 4.04(1H, m, OCH), δ 1.75(2H, t, Methylene) δ 1.38 (6H, d, Methyl).

EXAMPLE 5:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-6-pentadecyl-benzamide (CTPB):

25 2-ethoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted

into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo.

5 The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 554.6 (M+1), 359.6, 175.1, 149.1, 107.1, 91, 55.1, X-ray data also included

EXAMPLE 6:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-ethoxy-6-pentadecyl-benzamide:

2-ethoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-
10 trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-ethoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally
15 concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 545.5 (M+1), 359.6, 331.6, 213.3, 175.1, 149.1, 107.1, 91, 55.1

EXAMPLE 7:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-methoxy-6-pentadecyl-benzamide:

20 2-Methoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-methoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold
25 aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 540.4 (M+1), 345.6, 161.1, 149.1, 121.1, 91, 55.1

EXAMPLE 8:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-methoxy-6-pentadecyl-benzamide:

2-Methoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid
5 scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-methoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to
10 afford the desired product.

m/z: 531.6 (M+1), 345.7, 161, 149.2, 121.1, 91, 55.1

EXAMPLE 9:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-n-propoxy-6-pentadecyl-benzamide:

2-Propoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro
15 benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-n-propoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally
20 concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 568.8 (M+1), 526, 373, 331, 236, 147, 133, 107, 83, 55

EXAMPLE 10:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-isopropoxy-6-pentadecyl-benzamide:

25 2-Isopropoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-isopropoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was

extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

5 m/z: 568.8(M+1), 526, 373, 331, 236, 147, 133, 107, 83, 55

EXAMPLE 11:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-n-propoxy-6-pentadecyl-benzamide:

2-Propoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid
10 scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-n-propoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to
15 afford the desired product.

m/z: 560 (M+1), 517, 446, 373, 331, 147, 107, 91

EXAMPLE 12:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-isopropoxy-6-pentadecyl-benzamide:

2-Isopropoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid
20 scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-isopropoxy-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally
25 concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 560 (M+1), 517, 446, 373, 331, 147, 107, 91, 55.1

EXAMPLE 13:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-benzamide (CTB):

2-ethoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger
5 to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired
10 product.

m/z: 344.4 (M+1), 239, 149, 121, 102, 82, 57

EXAMPLE 14:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-ethoxy-benzamide:

2-ethoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-
15 trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-ethoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo.
20 The residue obtained was chromatographed over silica gel to afford the desired product.

¹H-NMR: δ 6.95-8.01 (7H, aromatic), δ 3.98 (2H, m, OCH₂), δ 1.33 (3H, t, Methyl)

EXAMPLE 15:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-methoxy-benzamide:

25 2-Methoxy-6-pentadecyl-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-methoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous

hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

¹H-NMR: δ6.95-7.40 (7H, aromatic), δ3.73(3H, s, OCH₃)

5 EXAMPLE 16:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-methoxy-benzamide:

2-Methoxy-6-pentadecyl-benzoyl chloride was condensed with 4-Amino-2-trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-methoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

¹H-NMR: δ6.95-8.01 (7H, aromatic), δ3.73(3H, OCH₃)

EXAMPLE 17:

N-(4-Chloro-3-trifluoromethyl-phenyl)-2-n-propoxy-benzamide:

2-Propoxy-benzoyl chloride was condensed with 5-amino-2-chloro benzenetrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield N-(4-Chloro-3-trifluoromethyl-phenyl)-2-n-propoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

¹H-NMR: δ6.95-8.01(7H, aromatic), δ3.94(2H, t, OCH₂), δ1.75(2H, m, Methylene), δ0.96(3H, t, Methyl)

EXAMPLE 18:

N-(4-Cyano-3-trifluoromethyl-phenyl)-2-Isopropoxy-benzamide:

2-Isopropoxy-benzoyl chloride was condensed with 4-Amino-2-trifluoromethyl benzonitrile in dichloromethane in presence of triethylamine as acid scavenger to
5 yield N-(4-Cyano-3-trifluoromethyl-phenyl)-2-Isopropoxy-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired
10 product.

¹H-NMR: δ6.95-8.01 (7H, aromatic), δ4.04 (1H, m, OCH), δ1.38 (6H, d, Methyl)

EXAMPLE 19:

2-Ethoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide:

2-ethoxy-6-pentadecyl-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in dichloromethane in presence of triethylamine as acid
15 scavenger to yield 2-Ethoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally
20 concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 565 (because of NO₂), 359, 175, 149, 136, 107, 77

EXAMPLE 20:

2-Methoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide:

25 2-Methoxy-6-pentadecyl-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield 2-Methoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with

cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

m/z: 551(because of NO₂),345, 161,149,121,91

5 **EXAMPLE 21:**

2-Propoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide:

2-Propoxy-6-pentadecyl-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in dichloromethane in presence of triethylamine as acid scavenger to yield 2-Propoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-
10 benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to afford the desired product.

15 m/z: 580(because of NO₂),537,373,331,189,161,147,91

EXAMPLE 22:

2-Isopropoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide:

2-Isopropoxy-6-pentadecyl-benzoyl chloride was condensed with 5-Amino-2-nitrobenzotrifluoride in dichloromethane in presence of triethylamine as acid
20 scavenger to yield 2-Isopropoxy-N-(4-nitro-3-trifluoromethyl-phenyl)-6-pentadecyl-benzamide. The reaction mixture was then concentrated in vacuo and the residue was extracted into ethyl acetate. The ethyl acetate layer was washed with water and with cold aqueous hydrochloric acid, then dried over sodium sulphate and finally concentrated in vacuo. The residue obtained was chromatographed over silica gel to
25 afford the desired product.

m/z: 580(because of NO₂),537,373,331,189,161,147,107,91

EXAMPLE 23:**Anacardic acid**

Isolation of Anacardic Acid from CNSL. Commercially available solvent-extracted CNSL (1Kg) was dissolved in 5% aqueous methanol (6.2L), and calcium hydroxide (510 g) was added in portions under stirring. After complete addition of calcium hydroxide, the temperature of the reaction mixture was raised to 54 °C and stirring was continued for 4h. The supernatant solution was monitored by TLC for the absence of anacardic acid. After completion of the reaction, the precipitated calcium anacardate was filtered and washed thoroughly with methanol (2.5L), and the cake was dried under vacuum at 45-54 °C for 3h (dry weight 1.08 Kg). The filtrate was preserved for subsequent isolation of cardol and cardanol. Calcium anacardate (1.08Kg) was suspended in distilled water (4.5 L) and 11 M HCl (600 mL) was added and stirred for 1.5 h. The resultant solution was extracted with ethyl acetate (2 × 2 L). The combined organic layer was washed with distilled water (2 × 2L), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to yield 600 g of mixture of anacardic acid (monoene, diene, and triene).

Preparation of Saturated Anacardic Acid. To a solution of an ene mixture of anacardic acid (500 g, 1300 mmol) in methanol (2.0 L) was added 5% palladium-carbon (25 g), and then hydrogen gas was passed through the solution at 2.5 kg/cm² until consumption of hydrogen gas ceased (4-5 h). The reaction mass was filtered over a Celite bed and washed with methanol (100 mL). The filtrate was concentrated under reduced pressure to give an off-white solid (477 g), which on recrystallization from hexane yielded 450g.

EXAMPLE 24:**Anacardic alcohol**

To a stirred solution of Ethyl 2-ethoxy-6-pentadecylbenzoate was reduced to alcohol using lithium aluminum hydride.

m/z: 334.9(M+1)317,163,147,133,121,107,91,69,55

EXAMPLE 25:

Anacardic aldehyde

Anacardic alcohol was oxidized to corresponding anacardic aldehyde using pyridinium chloro chromate.

5 m/z: 334.9(M+1)317,163,147,133,121,107,91,69,55

EXAMPLE 26:

2-Ethoxy-6-pentadecyl-benzoic acid:

To a stirred solution of Ethyl 2-Ethoxy-6-pentadecylbenzoate (20 g, 240 mmol) in dimethyl sulfoxide (80 mL) was added potassium *tert*-butoxide (20 g, 890 mmol) in
10 portions. The solution was heated to 68 °C on a water bath for 3 h, the reaction was monitored by TLC using a hexane-ethyl acetate (9:1) solvent system. The reaction mass was cooled to 15°C, poured into ice water, and then acidified with 10% dilute hydrochloric acid. The precipitated solid was filtered and washed thoroughly with distilled water, and the crude mass was recrystallized in petroleum ether (50 mL) to
15 yield an off-white solid of 2-Ethoxy-6-pentadecylbenzoic Acid (15 g, 80%).

¹H-NMR:δ6.80-7.44 (3H,aromatic), δ3.98(2H,m,OCH₂), δ2.55(2H,m,Ar-CH₂),1.29-1.63 (26H,m,methylene), δ1.33(3H,t,methyl next to OCH₂), δ0.96 (3H,t,Methyl)

EXAMPLE 27:

Cardanol:

20 Commercially available solvent-extracted CNSL (1Kg) was dissolved in 5% aqueous methanol (6.2L), and calcium hydroxide (510 g) was added in portions under stirring. After complete addition of calcium hydroxide, the temperature of the reaction mixture was raised to 54 °C and stirring was continued for 4h. The supernatant solution was monitored by TLC for the absence of anacardic acid. After completion of the reaction,
25 the precipitated calcium anacardate was filtered and washed thoroughly with methanol (2.5L). The methanol layer was collected and concentrated in vacuo. The

concentrate was further purified on silica gel (100-200 mesh) by column chromatography by using increasing amounts ethyl acetate in hexane to fractionate cardanol.

m/z: 304.7(M+1)149,121,107,95,71,57

5 **EXAMPLE 28:**

Cardol:

Commercially available solvent-extracted CNSL (1Kg) was dissolved in 5% aqueous methanol (6.2L), and calcium hydroxide (510 g) was added in portions under stirring. After complete addition of calcium hydroxide, the temperature of the reaction mixture
10 was raised to 54 °C and stirring was continued for 4h. The supernatant solution was monitored by TLC for the absence of anacardic acid. After completion of the reaction, the precipitated calcium anacardate was filtered and washed thoroughly with methanol (2.5L). The methanol layer was collected and concentrated in vacuo. The concentrate was further purified on silica gel (100-200 mesh) by column
15 chromatography by using increasing amounts ethyl acetate in hexane to fractionate cardol.

m/z: 321(M+1),149,123,107,95,69,55

Results:

We have identified a small molecule compound, anacardic acid, from Cashew Nut
20 Shell Liquid, know to have anti-tumor activity, which inhibits HAT activity of p300 and PCAF (Figure 1).

Surprisingly, the amide derivatives (Figure 2 and 3) of the same compound show an enhancement of p300 HAT activity with human core histones (Figure 4). These compounds are found to be specific for p300 since even at high concentration it
25 cannot affect the HAT activity of PCAF (Figure 5).

The inhibitor anacardic acid also inhibits p300 HAT activity dependent transcription from the chromatin template but not DNA transcription. These results indicate the HAT specific activity of anacardic acid. As expected, the amide derivatives enhance

HAT-dependent chromatin transcription whereas transcription from the DNA template remained unaffected.

- Most of the analogs of the amide compounds showed similar activity with regards to activation of histone acetylation, except one of the CN- derivatives, predominantly enhanced the acetylation of histone H3.

Brief Description of the Figures:

- Figure 1 .Different unsaturated anacardic acids present in CNSL.
- Figure 2 . General formulae of the compounds claimed for the patent.
- Figure 3. Effect of increasing concentration of Anacardic acid on the histone acetyltransferase activity of p300 and PCAF.
- Figure 4. X-ray crystal structure of N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-6-pentadecyl-benzamide (CTPB)
- Figure 5. X-ray crystal structure of N-(4-Chloro-3-trifluoromethyl-phenyl)-2-ethoxy-benzamide (CTB)
- Figure 6. Effect of increasing concentrations of CTPB and CTB on the HAT activity of p300.
- Figure 7. Specificity of CTPB to activate the histone transferase activity of p300 and not that of PCAF.

References

1. Roth, S. Y., Denu, J. M., and Allis, C. D. (2001) *Annu. Rev. Biochem.* **70**, 81–120
2. Sterner, D. E., and Berger, S. L. (2000) *Microbiol. Mol. Biol. Rev.* **64**, 2, 435–459
3. Redner, R. L., Wang, J., and Liu, J. M. (1999) *Blood* **94**, 2, 417–428
4. Marks, P. A., Rifkind, R. A., Richon, V. M., Breslow, R., Miller, T., and Kelly, W. K. (2001) *Nat. Rev. Cancer* **1**, 194–202
5. Giles, R. H., Peters, D. J., and Breuning, M. H. (1998) *Trends Genet.* **14**, 178–183

6. Murata, T., Kurokawa, R., Krones, A., Tatsumi, K., Ishii, M., Taki, T., Masuno, M., Ohashi, H., Yanagisawa, M., Rosenfeld, M. G., Glass, C. K., and Hayashi Y. (2001) *Hum. Mol. Genet.* **10**, 1071–1076
- 5 7. Ait-Si-Ali, S., Ramirez, S., Barre, F. X., Dkhissi, F., Magnaghi-Jaulin, L., Girault, J. A., Robin, P., Knibiehler, M., Pritchard, L. L., Ducommun, B., Trouche, D., and Harel-Bellan, A. (1998) *Nature*, **396**, 184–186
8. Kundu, T. K., Palhan, V., Wang, Z., An, W., Cole, P. A., and Roeder, R. G. (2000) *Mol. Cell* **6**, 551–561
9. Cullis, P. M., Wolfeuden, R., Cousens, L., and Alberts, B. M. (1982) *J. Biol. Chem.* **257**, 12165–12169
- 10 10. Lau, O. D., Kundu, T. K., Soccio, R. E., Ait-Si-Ali, S., Khalil, E. M., Vassilev, A., Wolfe, A. P., Nakatani, Y., Roeder, R. G., and Cole, P. A. (2000) *Mol. Cell* **5**, 589–595
11. Costanzo, A., Merlo, P., Pediconi, N., Fulco, M., Sartorelli, V., Cole, P. A., Fontemaggi, G., Fanciulli, M., Schiltz, L., Blandino, G., Balsano, C., and Levrero, M. (2002) *Mol. Cell* **9**, 175–186
- 15 12. Kundu, T. K., Wang, Z., and Roeder, R. G. (1999) *Mol. Cell. Biol.* **19**, 1605–1615
13. Wang, Z., and Roeder, R. G. (1997) *Genes Dev.* **11**, 1315–1326
- 20 14. Prashanth Kumar, B. R., Swaminathan, V., Banerjee, S., and Kundu, T. K. (2001) *J. Biol. Chem.* **276**, 16804–16809

CLAIMS

1. A compound of structural formula (I), as an activators of Histone acetyltransferases, *containing ring A derived from substituted benzoic acid moiety and ring B is substituted anilide* wherein:
 - 5 **R1** is H, Methyl, Ethyl, n-Propyl, Isopropyl, n-butyl, t-butyl, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;
 - R2** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl;
 - R3** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF_3 , CCl_3 , Cl_3 , F, Cl, I, NO_2 , CN;
 - 10 **R4** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF_3 , CCl_3 , Cl_3 , F, Cl, I, NO_2 , CN;
 - R5** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF_3 , CCl_3 , Cl_3 , F, Cl, I, NO_2 , ;
 - R6** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF_3 , CCl_3 , Cl_3 , F, Cl,
 - 15 I, NO_2 , CN; and
 - R7** is H, methyl, ethyl, n-propyl, isopropyl, n-butyl and t-butyl, CF_3 , CCl_3 , Cl_3 , F, Cl, I, NO_2 , CN.
2. A compound of structural formula (II) *for ring A of formula (I)* the accompanying drawings for inhibitor of Histone acetyltransferases, wherein:
 - 20 **R1** is H, CH_3 , Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy , O-Isopropoxy, n-butoxy, t-butoxy, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;
 - R2** is H, CH_3 , Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy , O-Isopropoxy, n-butoxy, t-butoxy, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;
 - R3** is H, CH_3 , Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy , O-
 - 25 Isopropoxy, n-butoxy, t-butoxy, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;
 - R4** is H, CH_3 , Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy , O-Isopropoxy, n-butoxy, t-butoxy, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;
 - R5** is H, CH_3 , Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy , O-Isopropoxy, n-butoxy, t-butoxy, C_8H_{18} , $C_{15}H_{26}$, $C_{15}H_{28}$, $C_{15}H_{30}$, $C_{15}H_{32}$;

30

R6 is H, CH₃, Hydroxyl, Carboxylic, O-Methoxy, O-Ethoxy, n-Propoxy, O-Isopropoxy, n-butoxy, t-butoxy, C₈H₁₈, C₁₅H₂₆, C₁₅H₂₈, C₁₅H₃₀, C₁₅H₃₂.

3. A process of preparing compounds as described in formula I or formula II by known methods.

- 5 4. A method of treating a patient suffering from diseases due defects in gene regulation predominantly or at risk of, cancer, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I) to activate histone acetyltransferases or formula (II) to inhibit histone acetyltransferase or a pharmaceutically acceptable salt or solvate of these compounds.

10

15

20

25

1/6






FIGURE-1



Formula I



Formula II

FIGURE-2

2/6




FIGURE-3

4/6




FIGURE-6

5/6




FIGURE -6




FIGURE-7

THIS PAGE BLANK (USPTO)